

A NEW CLASS OF RECONSTRUCTIBLE GRAPHS

TETSUYA HOSAKA

ABSTRACT. In this paper, we give a new class of reconstructible graphs.

1. INTRODUCTION

The purpose of this paper is to give a new class of reconstructible graphs.

For a graph G and $v \in V(G)$, let $N(v) = \{u \in V(G) \mid uv \in E(G)\}$ and $N[v] = N(v) \cup \{v\}$.

We prove the following theorem which is an extension of a result in [1].

Theorem 1.1. *Let G be a graph with $V(G) = \{v_1, \dots, v_k, \dots, v_n\}$ ($1 < k < n$). Suppose that*

- (1) $\bigcup_{i=1}^k N[v_i] = V(G)$,
- (2) $N[v_1] \cap \bigcup_{i=2}^k N[v_i] = \emptyset$,
- (3) $|d(v_i) - d(v_j)| \neq 1$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} - \{i\}$,
- (4) $\{v \in V(G) \mid d(v) = d(v_1)\} = \{v_1\}$, and
- (5) $\{v \in V(G) \mid d(v) = d(v_i)\} \subset \{v_2, \dots, v_k\}$ for any $i \in \{2, \dots, k\}$.

Then G is reconstructible.

2. PROOF OF THE THEOREM

Let G be a graph with $V(G) = \{v_1, \dots, v_k, \dots, v_n\}$ ($1 < k < n$). Suppose that

- (1) $\bigcup_{i=1}^k N[v_i] = V(G)$,
- (2) $N[v_1] \cap \bigcup_{i=2}^k N[v_i] = \emptyset$,

Date: July 23, 2004.

Key words and phrases. the reconstruction conjecture, reconstructible graphs.

Partly supported by the Grant-in-Aid for Scientific Research, The Ministry of Education, Culture, Sports, Science and Technology, Japan, (No. 15740029).

- (3) $|d(v_i) - d(v_j)| \neq 1$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, n\} - \{i\}$,
- (4) $\{v \in V(G) \mid d(v) = d(v_1)\} = \{v_1\}$, and
- (5) $\{v \in V(G) \mid d(v) = d(v_i)\} \subset \{v_2, \dots, v_k\}$ for any $i \in \{2, \dots, k\}$.

Let G' be a graph with $V(G') = \{v'_1, \dots, v'_n\}$ such that $G - v_j \cong G' - v'_j$ for any $j \in \{1, \dots, n\}$.

Here we note that $d(v_j) = d(v'_j)$ for any $j \in \{1, \dots, n\}$.

Let $f_j : G - v_j \rightarrow G' - v'_j$ be an isomorphism for each $j \in \{1, \dots, n\}$.

Lemma 2.1. $N[v'_1] \cap \bigcup_{i=2}^k N[v'_i] = \emptyset$.

Proof. Suppose that there exists $j \in \{1, \dots, n\}$ such that $v'_j \in N[v'_1] \cap \bigcup_{i=2}^k N[v'_i]$. Then $v'_j \in N[v'_1] \cap N[v'_{i_0}]$ for some $i_0 \in \{2, \dots, k\}$.

If $j = 1$ or $j = i_0$ then $v'_{i_0} \in N[v'_1]$. Here we consider an isomorphism $f_{i_0} : G - v_{i_0} \rightarrow G' - v'_{i_0}$. Since $|d(v_1) - d(v_m)| > 1$ for any $m \in \{2, \dots, n\}$ by (3) and (4), $f_{i_0}(v_1) = v'_1$ because $d(v_1) = d(v'_1)$. Let $d(v_1; G - v_{i_0})$ be the degree of v_1 in $G - v_{i_0}$. Then

$$\begin{aligned} d(v_1; G - v_{i_0}) &= d(f_{i_0}(v_1); G' - v'_{i_0}) = d(v'_1; G' - v'_{i_0}) \\ &= d(v'_1) - 1 = d(v_1) - 1. \end{aligned}$$

Hence $v_{i_0} \in N[v_1]$. This contradicts (2).

Suppose that $j \notin \{1, i_0\}$. We consider an isomorphism $f_j : G - v_j \rightarrow G' - v'_j$. Then $f_j(v_1) = v'_1$ by (3) and (4), and $f_j(v_{i_1}) = v'_{i_0}$ for some $i_1 \in \{2, \dots, k\}$ such that $d(v_{i_1}) = d(v_{i_0})$ by (3) and (5). Hence

$$\begin{aligned} d(v_1; G - v_j) &= d(f_j(v_1); G' - v'_j) = d(v'_1; G' - v'_j) \\ &= d(v'_1) - 1 = d(v_1) - 1, \end{aligned}$$

and

$$\begin{aligned} d(v_{i_1}; G - v_j) &= d(f_j(v_{i_1}); G' - v'_j) = d(v'_{i_0}; G' - v'_j) \\ &= d(v'_{i_0}) - 1 = d(v_{i_1}) - 1. \end{aligned}$$

This means that $v_j \in N[v_1] \cap N[v_{i_1}]$ which contradicts (2).

Thus $N[v'_1] \cap \bigcup_{i=2}^k N[v'_i] = \emptyset$. □

We consider an isomorphism $f_1 : G - v_1 \rightarrow G' - v'_1$. By (1) and (2),

$$f_1(V(G) - N[v_1]) = f_1\left(\bigcup_{i=2}^k N[v_i]\right).$$

Since $f_1(\{v_2, \dots, v_k\}) = \{v'_2, \dots, v'_k\}$ by (3) and (5),

$$f_1\left(\bigcup_{i=2}^k N[v_i]\right) = \bigcup_{i=2}^k N[v'_i].$$

By Lemma 2.1,

$$\bigcup_{i=2}^k N[v'_i] \subset V(G') - N[v'_1].$$

Thus

$$f_1(V(G) - N[v_1]) \subset V(G') - N[v'_1].$$

Here

$$\begin{aligned} |f_1(V(G) - N[v_1])| &= n - d(v_1) - 1 \\ &= n - d(v'_1) - 1 = |V(G') - N[v'_1]|. \end{aligned}$$

Hence $f_1(V(G) - N[v_1]) = V(G') - N[v'_1]$. This implies that $f_1(N(v_1)) = N(v'_1)$. Thus the map $f : G \rightarrow G'$ defined by $f(v_1) = v'_1$ and $f(v_j) = f_1(v_j)$ for each $j \in \{2, \dots, n\}$ is an isomorphism. Therefore G is reconstructible.

REFERENCES

- [1] T. Hosaka, *A class of reconstructible graphs*, preprint.
- [2] F.S. Mulla, *A class of graphs for which the Ulam conjecture holds*, Discrete Math. **22** (1978), 197–198.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA,
321-8505, JAPAN

E-mail address: hosaka@cc.utsunomiya-u.ac.jp